


HPC & SASSIE modelling of detergent micelles



Chris Lorenz




Biophysics & Soft Matter Group

Theory & Simulation of Condensed Matter Group

Physics Department

King's College London


Large q comparison to neutrons





Dan Allen – PhD student
Self-assembly of soft matter
Drug encapsulation

Prof. Jayne Lawrence, Dr. David Barlow & Yussif Saaka (KCL IPS): Neutron scattering & other biophysical experimental techniques




Introduction

- Self-assembly of soft nanomaterials proves to be very challenging for molecular simulations
 - Time scale of self-assembly is on the order of 10^{-4} s
 - Typical concentrations of experimental interest are on the order of 10^{-5} – 10^{-3} M, with aggregate sizes on the order of 100 molecules
 - Properties of interest for micelles are often concentration dependent
 - Therefore in explicit solvent systems that results in systems on the order of 10^5 - 10^7 atoms


Background

- Sodium dodecyl sulfate (SDS) is a commonly used surfactant in the pharmaceutical industry as a wetting and solubilising agent.
 - Antimalarial drug (Krishna et al. *J. Pharm. Sci.* **78**, 574-576 (1989))
 - Ibuprofen (Rangel-Yagui et al. *J. Pharm. Sci.* **41**, 237-246 (1996))
 - Anti-cancer drug (Enache et al. *Int. J. Pharm.* **390**, 100-106 (2010))



Background

- Testosterone propionate
 - Steroid used in hormone replacement therapy
 - Little is known regarding interactions with solubilising agents
 - Contains both **hydrophilic** (O1, O2 & O3) and **hydrophobic** regions which makes it difficult to predict where it will reside within micelle




Simulation set-up

Pre-assembled SDS micelle with 80 molecules

CHARMM forcefield w/ TIP3P for water

Concentration of SDS is 3 g/mol (same as exp't.)

Allen et al. *J. Phys. Chem. B* **118**, 13192-13201 (2014)





Simulation set-up

Pre-assembled SDS micelle with 80 molecules


Equilibrated SDS micelle (76 molecules) before surrounding with 13 TSTP molecules

CHARMM forcefield w/ TIP3P for water

Concentration of SDS is 3 g/mol (same as exp't.)




Resulting aggregates





	Blank SDS micelle	SDS micelle with TSTP
Eccentricity	0.11 ± 0.07	0.16 ± 0.08
I_x/I_y	1.01 ± 0.17	1.03 ± 0.18
I_x/I_z	1.04 ± 0.20	1.18 ± 0.29
I_y/I_z	1.04 ± 0.18	1.15 ± 0.28
Max l_x (Å)	48.06 ± 4.98	48.22 ± 5.81
Max l_y (Å)	48.21 ± 4.37	49.75 ± 5.67
Max l_z (Å)	49.18 ± 5.01	53.15 ± 6.06
SASA/molecule (Å ²)	169.9 ± 4.5	156.1 ± 4.2
Total SASA (Å ²)	13592.8 ± 358.2	13734.9 ± 371.6
Voronoi Surface Area/molecule (Å ²)	149.7 ± 2.6	136.9 ± 2.2
Total Voronoi Surface Area (Å ²)	11972.1 ± 206.6	12181.2 ± 193.5

Where are ions & water located?




Where is drug located?



Found drug was just under head group — useful for fitting neutron data


Drug moves on surface



Application of SASSIE

- Tried to use SASSIE to generate SANS profile from simulated structure of SDS micelle
- Read in CHARMM topology file for SDS
- Converted LAMMPS trajectory to dcd files
- Used all-atom SAScalc (beta version)

Application of SASSIE



At large Q the SANS profiles agree well

At small Q , not so much ... repulsive behaviour between micelles is not modelled by SASSIE

Further thoughts

- Simulations need aggregation number from experiment.
- Simulations can provide lots of relevant structural properties that can be used to gain molecular scale understanding of micelles and also help to fit neutron scattering data
- Still some way to go to be able to directly compare SANS and MD data

The logo for the Engineering and Physical Sciences Research Council (EPSRC). It features the acronym "EPSRC" in a bold, purple, sans-serif font. The letters are centered between two horizontal teal lines.

Engineering and Physical Sciences
Research Council